

Corn and Small Grain Silage Tests in Tennessee

2011

Fred L. Allen, Coordinator, Agronomic Crop Variety Testing & Demonstrations

Richard Johnson, Research Associate, Agronomic Crop Variety Testing & Demonstrations

Virginia Sykes, Graduate Research Assistant

**Agronomic Crop Variety Testing and Demonstrations
Department of Plant Sciences
Institute of Agriculture
University of Tennessee
Knoxville**

•Telephone: (865)974-8821 • FAX: (865)974-1947 •email: allenf@utk.edu

Variety test results are posted on UT's website at:

<http://varietytrials.tennessee.edu/>
and
www.utcrops.com

Acknowledgments

This research was funded by the Tennessee Agricultural Experiment Station and UT Extension with partial funding from participating companies.

We gratefully acknowledge the assistance of the following individuals in conducting these experiments:

Department of Plant Sciences

Dr. Dennis West, Professor and Grains Breeder

Mr. David Kincer, Research Associate

Research and Education Centers:

East Tennessee, Knoxville

Dr. Robert Simpson, Superintendent

Mr. Bobby McKee, Sr. Farm Crew Leader

Mr. Lee Ellis, Research Assistant

Plateau, Crossville

Mr. Walt Hitch, Superintendent

Mr. Greg Blaylock, Light Farm Equipment Operator

Mr. Sam Simmons, Light Farm Equipment Operator

Highland Rim, Springfield

Dr. Barry Sims, Superintendent

Mr. Brad S. Fisher, Research Associate

Middle Tennessee, Spring Hill

Dr. Kevin Thompson, Superintendent

Mr. Roy Thompson, Research Associate

County Standard Corn Silage Tests

County

Blount

Washington

Producer

Mac Pate

David Saylor

Agent

John Wilson

John Hamrick

Table of Contents

Experimental Procedures	3
Interpretation of Data	4
Research and Education Center Information	4
2011 Corn Hybrid Silage Yield	5
2011 Corn Hybrid Silage Agronomic Data	6
2011 Corn Hybrid Silage Quality Data	7
2 Year Corn Hybrid Silage Data	9
3 Year Corn Hybrid Silage Data	11
County Standard Tests	13
2011 Wheat Silage Yield & Agronomic Data	15
2011 Wheat Silage Quality Data	18
2 Year Wheat Silage Data	20
2011 Oat Silage Yield & Agronomic Data	23
2011 Oat Silage Quality Data	24
2 Year Oat Silage Data	25
Corn Hybrid Characteristics	26
Seed Company Contact Information	27

CORN SILAGE YIELD TESTS

2011

Experimental Procedures

Research and Education Center Tests: Twenty-two corn hybrids were evaluated for silage yield and quality in 2011. The tests were conducted at the East Tennessee (Knoxville), Plateau (Crossville), Highland Rim (Springfield), and Middle Tennessee (Spring Hill), Research and Education Centers (REC). The plots at all locations consisted of two rows, planted 30 inches apart, 30 feet in length and were replicated three times. Yields presented were adjusted to both dry weight and 65% moisture. The plant populations as well as the planting and harvesting dates are given in Table 1. Plots were harvested by commercial silage harvesters. A sub-sample from each plot of approximately 3 lbs was taken for analysis. Fresh weight and dried weight was recorded on each sample for determination of moisture at harvest. The samples were then ground and analyzed for nutrient content. Silage quality analyses were provided by Cumberland Valley Analytical Services, Inc., Hagerstown, MD. Milk per ton and milk per acre calculations were performed using the University of Wisconsin Milk2006 program.

County Standard Tests: The County Standard Corn Silage Tests were conducted in Blount and Washington counties in Tennessee. Each hybrid was evaluated in a large strip-plot at each location, thus **each county test was considered as one replication of the test** in calculating the overall average yield and in conducting the statistical analysis to determine significant differences. At each location, plots were planted, sprayed, fertilized, and harvested with the equipment used in the cooperating producer's farming operation. The width and length of strip-plots were different in each county; however, within a location in a county, the strips were trimmed on the ends so that the lengths were the same for each variety, or if the lengths were different then the harvested length was measured for each variety and appropriate harvested area adjustments were made to determine the yield per acre.

Growing Season: The 2011 growing season was characterized by a wet spring resulting in some flooded fields followed by hotter and dryer than normal conditions. Hot dry conditions in July and August advanced harvest slightly ahead of the normal pace.

Interpretation of Data:

The tables on the following pages have been prepared with the entries listed in order of performance, the highest-yielding entry being listed first. At the bottom of the tables, **LSD** values stand for **Least Significant Difference**. The mean yields of any two varieties being compared must differ by at least the amount shown to be considered different in yielding ability at the 5% level of probability of significance. For example, given that the LSD for a test is 1.3 tons/a and the mean yield of Hybrid A was 9.3 tons/a and the mean yield of Hybrid B was 8.2 tons/a, then the two hybrids are not statistically different in yield because the difference of 1.1 tons/a is less than the minimum of 1.3 tons/a required for them to be significant. Similarly, if the average yield of Hybrid C was 10.6 tons/a then it is significantly higher yielding than both Hybrid B ($10.6 - 8.2 = 2.4$ tons/a > LSD of 1.3) and Hybrid A ($10.6 - 9.3 = 1.3$ tons/a = LSD of 1.3).

Also, the **coefficient of variation (C.V.)** values are shown at the bottom of each table. This value is a measure of the error variability found within each experiment. It is the percentage that the square root of error variance is of the overall test mean yield at that location. For example, a C.V. of 10% indicates that the size of the error variation is about 10% of the size of the test mean. Similarly, a C.V. of 30% indicates that the size of the error variation is nearly one-third as large as the test mean. A goal in conducting each yield test is to keep the C.V. as low as possible, preferably below 20 percent.

Table 1. Location information from Research and Education Centers where the corn silage variety tests were conducted in 2011.

Research and Education Center	Location	Planting Date	Harvest Date	Plant Population	Soil Type
East Tennessee	Knoxville	4/19/11	8/16/11	28,624	Sequatchie Silt Loam
Plateau	Crossville	5/11/11	8/24/11	30,492	Lilly Silt Loam
Middle Tennessee	Spring Hill	5/2/11	8/5/11	24,684	Maury Silt Loam
Highland Rim	Springfield	4/19/11	8/16/11	27,588	Stasser Silt Loam

Table 2. Mean yields † of 22 corn hybrids evaluated for silage at four locations in Tennessee during 2011.

Brand	Hybrid §	Dry Weight	65% Moisture	Dry Weight			
		Avg. Yield ± Std Err. (n=4)	Avg. Yield ± Std Err. (n=4)	Knoxville	Crossville	Spring Hill	Springfield
		tons/a					
Mycogen	TMF2H918 (RR/LL/HX)	8.1 ± 0.4	23.2 ± 1.2	10.7	6.3	4.8	10.7
Augusta	A5462GT3000A (RR/LL/CB/RW)	8.0 ± 0.4	22.9 ± 1.2	8.7	8.4	5.9	9.2
Augusta	A5461GTCBLLA	8.0 ± 0.5	22.9 ± 1.3	10.6	6.8	5.6	9.1
Mycogen	TMF2L872 (RR/LL/HX1)	8.0 ± 0.5	22.9 ± 1.3	10.1	7.0	4.9	10.0
Augusta	A5464GTCBLL	7.9 ± 0.5	22.6 ± 1.3	10.1	7.1	5.0	9.5
Augusta	A5560VT3	7.8 ± 0.5	22.4 ± 1.4	9.2	6.6	6.4	9.2
DeKalb	DKC67-21 GENVT3P	7.8 ± 0.5	22.2 ± 1.3	9.3	8.1	3.9	9.8
Augusta	A7664VT3	7.8 ± 0.5	22.2 ± 1.3	10.4	7.1	5.2	8.4
Wyffels	W8681 (VT3Pro)	7.7 ± 0.4	22.0 ± 1.2	8.9	7.2	4.9	9.7
Wyffels	W8437 (VT3Pro)	7.6 ± 0.5	21.8 ± 1.4	8.4	6.1	5.7	10.3
Croplan	8756VT3	7.6 ± 0.5	21.6 ± 1.4	10.7	6.0	4.8	8.7
Croplan	8505VT3P	7.5 ± 0.4	21.4 ± 1.2	10.6	5.4	5.8	8.2
Croplan	8221VT3	7.5 ± 0.4	21.3 ± 1.2	10.8	5.6	5.6	7.8
Croplan	9009 RH	7.4 ± 0.4	21.1 ± 1.2	11.5	5.1	4.8	8.2
Croplan	7131VT3	7.4 ± 0.4	21.0 ± 1.2	8.3	6.2	5.4	9.6
Augusta	A6867CBLL	7.3 ± 0.4	20.8 ± 1.2	8.9	6.0	5.2	9.0
Augusta	A6867GTCBLLA	7.2 ± 0.5	20.7 ± 1.3	8.4	6.3	5.1	9.1
Augusta	A5337EVT3	7.1 ± 0.4	20.4 ± 1.2	9.7	5.9	5.4	7.6
Augusta	A6166GT3000A (RR/LL/CB/RW)	7.0 ± 0.4	20.1 ± 1.2	11.0	4.2	5.1	7.8
Masters Choice	MCT-6583 (RR/LL/CB/RW)	6.7 ± 0.4	19.0 ± 1.2	9.6	5.8	4.5	6.7
Croplan	6286 SS (RR2/LL/CB/RW)	6.4 ± 0.5	18.2 ± 1.3	8.5	4.9	4.6	7.4
Masters Choice	MC-590	6.3 ± 0.5	18.0 ± 1.3	7.7	6.6	3.9	7.0
Avg. (tons/a)		7.5	21.3	9.6	6.2	5.1	8.8
L.S.D._{.05} (tons/a)		1.2	3.4	1.7	3.3	1.2	2.9
C.V. (%)		19.3	19.3	10.2	30.8	13.5	20.1

† all silage yields are adjusted to dry weight basis unless otherwise indicated.

YG, YGCB, Bt, HX, CB = contains a *Bacillus thuringiensis* gene for insect resistance

LL = contains a gene for tolerance to glufosinate

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

YGRW, RW, CRW = contains a gene for rootworm resistance

RR, RR2, R, GT, R2 = contains a gene for tolerance to glyphosate

CL = contains a gene for tolerance to Imidazolinone class herbicides

Table 3. Mean yields † and agronomic characteristics of 22 corn hybrids evaluated for silage at four locations in Tennessee during 2011.

Brand	Hybrid §	Dry Weight	65% Moisture	Moisture at harvest (n=4)	Lodging (n=4)	Plant Height (n=4)	Ear Height (n=3)
		Avg. Yield ± Std Err. (n=4)	Avg. Yield ± Std Err. (n=4)				
		tons/a	tons/a	%	%	inches	inches
Mycogen	TMF2H918 (RR/LL/HX)	8.1 ± 0.4	23.2 ± 1.2	58.3	0	108	40
Augusta	A5462GT3000A (RR/LL/CB/RW)	8.0 ± 0.4	22.9 ± 1.2	56.6	0	108	37
Augusta	A5461GTCBLLA	8.0 ± 0.5	22.9 ± 1.3	53.9	0	104	40
Mycogen	TMF2L872 (RR/LL/HX1)	8.0 ± 0.5	22.9 ± 1.3	55.7	0	104	38
Augusta	A5464GTCBLL	7.9 ± 0.5	22.6 ± 1.3	54.1	0	101	37
Augusta	A5560VT3	7.8 ± 0.5	22.4 ± 1.4	53.6	0	102	33
DeKalb	DKC67-21 GENVT3P	7.8 ± 0.5	22.2 ± 1.3	54.8	0	100	36
Augusta	A7664VT3	7.8 ± 0.5	22.2 ± 1.3	56.0	0	102	31
Wyffels	W8681 (VT3Pro)	7.7 ± 0.4	22.0 ± 1.2	55.8	0	97	33
Wyffels	W8437 (VT3Pro)	7.6 ± 0.5	21.8 ± 1.4	54.4	0	94	30
Croplan	8756VT3	7.6 ± 0.5	21.6 ± 1.4	57.8	0	108	38
Croplan	8505VT3P	7.5 ± 0.4	21.4 ± 1.2	56.9	0	108	41
Croplan	8221VT3	7.5 ± 0.4	21.3 ± 1.2	55.3	0	110	44
Croplan	9009 RH	7.4 ± 0.4	21.1 ± 1.2	56.3	0	111	42
Croplan	7131VT3	7.4 ± 0.4	21.0 ± 1.2	55.2	0	98	30
Augusta	A6867CBLL	7.3 ± 0.4	20.8 ± 1.2	54.8	0	109	38
Augusta	A6867GTCBLLA	7.2 ± 0.5	20.7 ± 1.3	55.8	0	106	38
Augusta	A5337EVT3	7.1 ± 0.4	20.4 ± 1.2	56.2	0	108	35
Augusta	A6166GT3000A (RR/LL/CB/RW)	7.0 ± 0.4	20.1 ± 1.2	56.9	0	106	41
Masters Choice	MCT-6583 (RR/LL/CB/RW)	6.7 ± 0.4	19.0 ± 1.2	58.2	0	106	37
Croplan	6286 SS (RR2/LL/CB/RW)	6.4 ± 0.5	18.2 ± 1.3	56.1	0	102	33
Masters Choice	MC-590	6.3 ± 0.5	18.0 ± 1.3	57.5	0	101	33
Average		7.5	21.3	55.9	0	104	37

† all silage yields are adjusted to dry weight basis unless otherwise indicated.

YG, YGCB, CB, Bt, HX = contains a *Bacillus thuringiensis* gene for insect resistance

LL = contains a gene for tolerance to glufosinate

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

YGRW, RW = contains a gene for rootworm resistance

RR, R, RR2, R2, GT = contains a gene for tolerance to glyphosate

CL = contains a gene for tolerance to Imidazolinone class herbicides

Table 4. Mean yields † and feed quality characteristics of 22 corn hybrids evaluated for silage at four locations in Tennessee during 2011.

Brand	Hybrid §	Dry Weight											
		Avg. Yield ± Std Err. (n=4)	Moisture at Harvest (n=4)	Crude Protein (n=4)	NDF 30h								
		tons/a	%	% dm	NDF (n=4)	IV Digest (n=4)	Starch (n=4)	ADF (n=4)	TDN (n=4)	NEL (n=4)	Milk/ton‡ (n=4)	Milk/acre‡ (n=4)	
Mycogen	TMF2H918 (RR/LL/HX)	8.1 ± 0.4	58.3	7.6	42.7	54.2	32.1	25.9	72.4	0.76	3036	24589	
Augusta	A5462GT3000A (RR/LL/CB/RW)	8.0 ± 0.4	56.6	7.4	41.6	56.2	35.4	24.5	72.7	0.76	2951	23697	
Augusta	A5461GTCBLLA	8.0 ± 0.5	53.9	7.3	40.0	57.8	37.3	23.5	73.6	0.77	2957	23743	
Mycogen	TMF2L872 (RR/LL/HX1)	8.0 ± 0.5	55.7	7.5	42.8	57.5	33.1	25.5	72.5	0.76	2975	23798	
Augusta	A5464GTCBLL	7.9 ± 0.5	54.1	7.6	39.3	55.9	37.3	23.3	73.7	0.77	2933	23232	
Augusta	A5560VT3	7.8 ± 0.5	53.6	7.6	41.6	57.7	35.4	24.6	72.8	0.76	2939	23011	
DeKalb	DKC67-21 GENVT3P	7.8 ± 0.5	54.8	7.6	40.4	57.8	35.5	24.0	73.2	0.77	2980	23124	
Augusta	A7664VT3	7.8 ± 0.5	56.0	7.8	42.1	56.1	33.4	24.9	72.6	0.76	2960	22973	
Wyffels	W8681 (VT3Pro)	7.7 ± 0.4	55.8	7.8	40.4	57.7	35.8	24.0	73.3	0.77	2988	22980	
Wyffels	W8437 (VT3Pro)	7.6 ± 0.5	54.4	7.8	36.9	58.4	39.8	21.6	74.9	0.79	3047	23252	
Croplan	8756VT3	7.6 ± 0.5	57.8	7.4	44.6	57.2	30.9	26.4	71.6	0.75	3014	22813	
Croplan	8505VT3P	7.5 ± 0.4	56.9	7.5	41.4	55.5	34.1	24.4	73.3	0.77	3014	22608	
Croplan	8221VT3	7.5 ± 0.4	55.3	7.6	41.7	54.5	35.5	24.9	72.6	0.76	2855	21271	
Croplan	9009 RH	7.4 ± 0.4	56.3	7.6	44.4	55.7	31.3	26.8	71.7	0.75	2936	21729	
Croplan	7131VT3	7.4 ± 0.4	55.2	7.9	39.2	57.2	36.8	22.9	73.9	0.78	2992	21990	
Augusta	A6867CBLL	7.3 ± 0.4	54.8	8.0	38.8	58.5	38.0	22.6	73.7	0.77	2980	21696	
Augusta	A6867GTCBLLA	7.2 ± 0.5	55.8	7.9	41.7	57.6	34.1	24.6	72.6	0.76	2965	21436	
Augusta	A5337EVT3	7.1 ± 0.4	56.2	7.6	41.2	56.1	35.3	24.4	72.8	0.76	2943	20984	
Augusta	A6166GT3000A (RR/LL/CB/RW)	7.0 ± 0.4	56.9	7.6	40.5	58.0	35.6	23.6	73.3	0.77	3051	21477	
Masters Choice	MCT-6583 (RR/LL/CB/RW)	6.7 ± 0.4	58.2	7.4	42.3	55.4	33.1	25.4	72.2	0.75	3026	20125	
Croplan	6286 SS (RR2/LL/CB/RW)	6.4 ± 0.5	56.1	7.9	39.3	56.0	37.2	22.8	73.9	0.77	2998	19095	
Masters Choice	MC-590	6.3 ± 0.5	57.5	7.5	40.8	57.4	35.0	24.4	73.1	0.77	3071	19317	

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

Bt, YG, YGCB, CB, HX = contains a *Bacillus thuringiensis* gene for insect resistance

YGRW, RW = contains a gene for rootworm resistance

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

RR, RR2 = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

CL = contains a gene for tolerance to Imidazolinone class herbicides

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

Table 5. Mean yields † and feed quality characteristics of 22 corn hybrids evaluated for silage at four locations in Tennessee during 2011, sorted by brand.

Brand	Hybrid §	Dry Weight											
		Avg. Yield ± Std Err. (n=4)	Moisture at Harvest (n=4)	Crude Protein (n=4)	NDF (n=4)	IV Digest (n=4)	Starch (n=4)	ADF (n=4)	TDN (n=4)	NEL (n=4)	Milk/ton‡ (n=4)	Milk/acre‡ (n=4)	
		tons/a	%	% dm	% dm	% of NDF	% dm	% dm	% dm	Mcals/lb	lbs/ton	lbs/acre	
Augusta	A5461GTCBLLA	8.0 ± 0.5	53.9	7.3	40.0	57.8	37.3	23.5	73.6	0.77	2957	23743	
Augusta	A5462GT3000A (RR/LL/CB/RW)	8.0 ± 0.4	56.6	7.4	41.6	56.2	35.4	24.5	72.7	0.76	2951	23697	
Augusta	A5464GTCBLL	7.9 ± 0.5	54.1	7.6	39.3	55.9	37.3	23.3	73.7	0.77	2933	23232	
Augusta	A5560VT3	7.8 ± 0.5	53.6	7.6	41.6	57.7	35.4	24.6	72.8	0.76	2939	23011	
Augusta	A7664VT3	7.8 ± 0.5	56.0	7.8	42.1	56.1	33.4	24.9	72.6	0.76	2960	22973	
Augusta	A6867CBLL	7.3 ± 0.4	54.8	8.0	38.8	58.5	38.0	22.6	73.7	0.77	2980	21696	
Augusta	A6867GTCBLLA	7.2 ± 0.5	55.8	7.9	41.7	57.6	34.1	24.6	72.6	0.76	2965	21436	
Augusta	A5337EVT3	7.1 ± 0.4	56.2	7.6	41.2	56.1	35.3	24.4	72.8	0.76	2943	20984	
Augusta	A6166GT3000A (RR/LL/CB/RW)	7.0 ± 0.4	56.9	7.6	40.5	58.0	35.6	23.6	73.3	0.77	3051	21477	
Croplan	8756VT3	7.6 ± 0.5	57.8	7.4	44.6	57.2	30.9	26.4	71.6	0.75	3014	22813	
Croplan	8505VT3P	7.5 ± 0.4	56.9	7.5	41.4	55.5	34.1	24.4	73.3	0.77	3014	22608	
Croplan	8221VT3	7.5 ± 0.4	55.3	7.6	41.7	54.5	35.5	24.9	72.6	0.76	2855	21271	
Croplan	9009 RH	7.4 ± 0.4	56.3	7.6	44.4	55.7	31.3	26.8	71.7	0.75	2936	21729	
Croplan	7131VT3	7.4 ± 0.4	55.2	7.9	39.2	57.2	36.8	22.9	73.9	0.78	2992	21990	
Croplan	6286 SS (RR2/LL/CB/RW)	6.4 ± 0.5	56.1	7.9	39.3	56.0	37.2	22.8	73.9	0.77	2998	19095	
DeKalb	DKC67-21 GENVT3P	7.8 ± 0.5	54.8	7.6	40.4	57.8	35.5	24.0	73.2	0.77	2980	23124	
Masters Choice	MCT-6583 (RR/LL/CB/RW)	6.7 ± 0.4	58.2	7.4	42.3	55.4	33.1	25.4	72.2	0.75	3026	20125	
Masters Choice	MC-590	6.3 ± 0.5	57.5	7.5	40.8	57.4	35.0	24.4	73.1	0.77	3071	19317	
Mycogen	TMF2H918 (RR/LL/HX)	8.1 ± 0.4	58.3	7.6	42.7	54.2	32.1	25.9	72.4	0.76	3036	24589	
Mycogen	TMF2L872 (RR/LL/HX1)	8.0 ± 0.5	55.7	7.5	42.8	57.5	33.1	25.5	72.5	0.76	2975	23798	
Wyffels	W8681 (VT3Pro)	7.7 ± 0.4	55.8	7.8	40.4	57.7	35.8	24.0	73.3	0.77	2988	22980	
Wyffels	W8437 (VT3Pro)	7.6 ± 0.5	54.4	7.8	36.9	58.4	39.8	21.6	74.9	0.79	3047	23252	

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

Bt, YG, YGCB, CB, HX = contains a *Bacillus thuringiensis* gene for insect resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

YGRW, RW = contains a gene for rootworm resistance

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

RR, RR2 = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

CL = contains a gene for tolerance to Imidazolinone class herbicides

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

Table 6. Mean yields † of 10 corn hybrids evaluated for silage in four environments for two years (2010-2011) in Tennessee.

Brand	Hybrid §	Dry Weight	65% Moisture	Dry Weight			
		Avg. Yield ± Std Err. (n=8)	Avg. Yield ± Std Err. (n=8)	Knoxville	Crossville	Spring Hill	Springfield
-----tons/a-----							
Mycogen	TMF2H918 (RR/LL/HX)	7.8 ± 0.3	22.1 ± 0.8	11.2	5.7	5.3	8.8
Croplan	8505VT3P	7.7 ± 0.3	21.9 ± 0.8	10.8	5.7	6.0	8.2
Augusta	A5461GTCBLLA	7.6 ± 0.3	21.8 ± 0.9	10.9	5.8	6.1	7.7
Croplan	9009 RH	7.5 ± 0.3	21.5 ± 0.8	11.8	5.5	5.1	7.7
Augusta	A5462GT3000A (RR/LL/CB/RW)	7.5 ± 0.3	21.4 ± 0.8	9.5	6.8	5.8	7.8
Augusta	A5337EVT3	7.4 ± 0.3	21.1 ± 0.8	10.8	5.4	5.5	7.8
Croplan	8756VT3	7.3 ± 0.3	21.0 ± 0.9	10.7	5.3	5.3	8.1
Wyffels	W8681 (VT3Pro)	7.3 ± 0.3	20.9 ± 0.8	9.8	6.3	5.2	8.1
Augusta	A6867CBLL	7.3 ± 0.3	20.8 ± 0.8	10.0	5.9	5.3	7.9
Croplan	8221VT3	7.2 ± 0.3	20.7 ± 0.8	11.2	5.3	5.5	7.1
Avg. (tons/a)		7.5	21.3	10.7	5.8	5.5	7.9
L.S.D._{.05} (tons/a)		1.0	2.8	1.5	2.4	1.2	2.8
C.V. (%)		18.1	18.1	9.4	26.9	14.7	22.4

Table 7. Mean yields † and agronomic characteristics of 10 corn hybrids evaluated for silage in four environments for two years (2010-2011) in Tennessee.

Brand	Hybrid §	Dry Weight	65% Moisture	Moisture	Plant	Ear
		Avg. Yield ± Std Err. (n=8)	Avg. Yield ± Std Err. (n=8)	at harvest (n=8)	Lodging (n=8)	Height (n=8)
tons/a tons/a % % inches inches						
Mycogen	TMF2H918 (RR/LL/HX)	7.8 ± 0.3	22.1 ± 0.8	55.9	0	109
Croplan	8505VT3P	7.7 ± 0.3	21.9 ± 0.8	55.9	0	107
Augusta	A5461GTCBLLA	7.6 ± 0.3	21.8 ± 0.9	53.1	0	108
Croplan	9009 RH	7.5 ± 0.3	21.5 ± 0.8	55.2	0	110
Augusta	A5462GT3000A (RR/LL/CB/RW)	7.5 ± 0.3	21.4 ± 0.8	54.5	0	111
Augusta	A5337EVT3	7.4 ± 0.3	21.1 ± 0.8	55.5	0	107
Croplan	8756VT3	7.3 ± 0.3	21.0 ± 0.9	56.0	0	105
Wyffels	W8681 (VT3Pro)	7.3 ± 0.3	20.9 ± 0.8	54.6	0	100
Augusta	A6867CBLL	7.3 ± 0.3	20.8 ± 0.8	53.2	0	107
Croplan	8221VT3	7.2 ± 0.3	20.7 ± 0.8	53.9	0	107

† all silage yields are adjusted to dry weight basis unless otherwise indicated.

YG, YGCB, CB, Bt, HX = contains a *Bacillus thuringiensis* gene for insect resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

RR, R, RR2, R2, GT = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

YGRW, RW = contains a gene for rootworm resistance

CL = contains a gene for tolerance to Imidazolinone class herbicides

Table 8. Mean yields † and feed quality characteristics of 10 corn hybrids evaluated for silage at four locations for 2 years (2010-2011) in Tennessee.

Brand	Hybrid §	Dry Weight											
		Avg. Yield ± Std Err. (n=8)	Moisture at Harvest (n=8)	Crude		NDF 30h				NEL (n=8)	Milk/ton‡ (n=8)	Milk/acre‡ (n=8)	
				Protein (n=8)	NDF (n=8)	IV Digest (n=8)	Starch (n=8)	ADF (n=8)	TDN (n=8)				
		tons/a	%	% dm	% dm	% of NDF	% dm	% dm	% dm	Mcals/lb	lbs/ton	lbs/acre	
Mycogen	TMF2H918 (RR/LL/HX)	7.8 ± 0.3	55.9	7.7	45.8	57.2	28.7	27.7	71.0	0.74	2983	23136	
Croplan	8505VT3P	7.7 ± 0.3	55.9	7.5	44.5	52.6	31.3	26.9	71.3	0.74	2819	21628	
Augusta	A5461GTCBLLA	7.6 ± 0.3	53.1	7.4	42.2	56.6	34.0	25.3	72.3	0.75	2900	22176	
Croplan	9009 RH	7.5 ± 0.3	55.2	7.4	46.4	53.6	28.7	28.3	70.4	0.73	2812	21141	
Augusta	A5462GT3000A (RR/LL/CB/RW)	7.5 ± 0.3	54.5	7.6	42.1	56.0	33.7	25.0	72.2	0.75	2926	21884	
Augusta	A5337EVT3	7.4 ± 0.3	55.5	7.7	41.9	55.7	33.1	25.2	72.2	0.75	2929	21580	
Croplan	8756VT3	7.3 ± 0.3	56.0	7.5	47.3	57.7	27.8	28.0	70.3	0.73	2930	21543	
Wyffels	W8681 (VT3Pro)	7.3 ± 0.3	54.6	8.1	42.6	57.0	33.0	25.4	72.1	0.75	2929	21476	
Augusta	A6867CBLL	7.3 ± 0.3	53.2	7.9	40.9	57.3	34.9	24.3	72.6	0.76	2932	21363	
Croplan	8221VT3	7.2 ± 0.3	53.9	7.6	43.6	54.1	32.4	26.4	71.3	0.74	2806	20337	

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

Bt, YG, YGCB, CB, HX = contains a *Bacillus thuringiensis* gene for insect resistance

YGRW, RW = contains a gene for rootworm resistance

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

RR, RR2 = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

CL = contains a gene for tolerance to Imidazolinone class herbicides

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

Table 9. Mean yields † of five corn hybrids evaluated for silage in four environments for three years (2009-2011) in Tennessee.

Brand	Hybrid §	Dry Weight	65% Moisture	Dry Weight			Springfield
		Avg. Yield ± Std Err. (n=12)	Avg. Yield ± Std Err. (n=12)	Knoxville	Crossville	Spring Hill	
		tons/a					
Mycogen	TMF2H918 (RR/LL/HX)	8.3 ± 0.2	23.6 ± 0.6	11.4	6.3	6.6	8.8
Croplan	9009 RH	8.2 ± 0.2	23.3 ± 0.6	12.1	6.4	6.6	7.6
Croplan	8756VT3	8.0 ± 0.2	22.9 ± 0.6	11.5	5.7	6.6	8.2
Wyffels	W8681 (VT3Pro)	7.9 ± 0.2	22.7 ± 0.6	10.4	6.4	6.4	8.6
Croplan	8221VT3	7.8 ± 0.2	22.4 ± 0.6	11.0	6.2	6.3	7.8
Avg. (tons/a)		8.0	23.0	11.3	6.2	6.5	8.2
L.S.D._{.05} (tons/a)		0.9	2.5	1.5	2.0	1.2	2.3
C.V. (%)		14.7	14.7	9.3	21.2	11.4	18.1

Table 10. Mean yields † and agronomic characteristics of five corn hybrids evaluated for silage in four environments for three years (2009-2011) in Tennessee.

Brand	Hybrid §	Dry Weight	65% Moisture				Ear Height (n=9)
		Avg. Yield ± Std Err. (n=12)	Avg. Yield ± Std Err. (n=12)	Moisture at harvest (n=12)	Lodging (n=11)	Plant Height (n=12)	
		tons/a	tons/a	%	%	inches	
Mycogen	TMF2H918 (RR/LL/HX)	8.3 ± 0.2	23.6 ± 0.6	58.4	0	111	48
Croplan	9009 RH	8.2 ± 0.2	23.3 ± 0.6	58.1	0	112	49
Croplan	8756VT3	8.0 ± 0.2	22.9 ± 0.6	57.2	0	106	45
Wyffels	W8681 (VT3Pro)	7.9 ± 0.2	22.7 ± 0.6	56.8	0	101	40
Croplan	8221VT3	7.8 ± 0.2	22.4 ± 0.6	56.1	0	107	49

Codes:

† all silage yields are adjusted to dry weight basis unless otherwise indicated.

YG, YGCB, CB, Bt, HX = contains a *Bacillus thuringiensis* gene for insect resistance

YGRW, RW = contains a gene for rootworm resistance

RR, R, RR2, R2, GT = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

CL = contains a gene for tolerance to Imidazolinone class herbicides

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

Table 11. Mean yields † and feed quality characteristics of five corn hybrids evaluated for silage at four locations for three years (2009-2011) in Tennessee.

Brand	Hybrid §	Dry Weight		Moisture		Crude		NDF 30h				NEL (n=12)	Milk/ton‡ (n=12)	Milk/acre‡ (n=12)
		Avg. Yield ± Std Err. (n=12)	tons/a	%	at Harvest (n=12)	Protein (n=12)	NDF (n=12)	IV Digest (n=12)	Starch (n=12)	ADF (n=12)	TDN (n=12)			
Mycogen	TMF2H918 (RR/LL/HX)	8.3 ± 0.2	58.4	7.7	46.2	56.1	27.2	28.4	71.1	0.74	3003	25469		
Croplan	9009 RH	8.2 ± 0.2	58.1	7.6	47.2	52.9	26.9	29.1	70.4	0.73	2889	23975		
Croplan	8756VT3	8.0 ± 0.2	57.2	7.5	46.5	57.6	28.5	27.7	70.9	0.74	2987	24234		
Wyffels	W8681 (VT3Pro)	7.9 ± 0.2	56.8	8.3	42.1	57.7	32.4	25.1	72.6	0.76	3048	24612		
Croplan	8221VT3	7.8 ± 0.2	56.1	7.6	43.0	54.3	32.1	26.2	71.9	0.75	2902	23317		

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

Bt, YG, YGCB, CB, HX = contains a *Bacillus thuringiensis* gene for insect resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

YGRW, RW = contains a gene for rootworm resistance

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

RR, RR2 = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

CL = contains a gene for tolerance to Imidazolinone class herbicides

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

COUNTY STANDARD TESTS

Table 12. Mean yields † of 20 corn hybrids evaluated for silage in two County Standard Tests in Tennessee during 2011.

Brand	Hybrid §	Dry Weight	65% Moisture			Moisture at harvest (n=2)	Lodging (n=1)	Plant Height (n=1)	Ear Height (n=1)
		Avg. Yield ± Std Err. (n=2)	Avg. Yield ± Std Err. (n=2)	----- Dry Weight -----	Blount (n=1)	Washington (n=1)			
		-----tons/a-----				%			
Augusta	A7664VT3	6.7 ± 0.4	19.1 ± 1.2	6.4	7.0	50.5	0	80	36
DeKalb	DKC67-21 GENVT3P	6.4 ± 0.4	18.4 ± 1.2	6.2	6.7	52.3	0	92	47
Croplan	7131VT3	6.3 ± 0.4	18.1 ± 1.2	4.9	7.7	59.3	0	76	36
Augusta	A5462GT3000A (RR/LL/CB/RW)	6.3 ± 0.4	17.9 ± 1.2	5.4	7.1	56.5	0	84	45
Croplan	8756VT3	6.1 ± 0.4	17.6 ± 1.2	4.9	7.4	63.1	0	88	46
Wyffels	W8681 (VT3Pro)	6.0 ± 0.4	17.3 ± 1.2	5.6	6.5	56.5	0	78	38
Croplan	8505VT3P	6.0 ± 0.4	17.0 ± 1.2	5.6	6.3	57.3	0	88	48
Croplan	6286 SS (RR2/LL/CB/RW)	6.0 ± 0.4	17.0 ± 1.2	4.5	7.4	60.0	0	84	45
Masters Choice	MCT-6583 (RR/LL/CB/RW)	5.9 ± 0.4	16.9 ± 1.2	5.1	6.7	59.1	0	90	48
Augusta	A5461GTCBLLA	5.9 ± 0.4	16.8 ± 1.2	4.8	7.0	58.1	0	96	46
Mycogen	TMF2L872 (RR/LL/HX1)	5.9 ± 0.4	16.8 ± 1.2	4.5	7.3	65.1	0	96	48
Augusta	A6867GTCBLLA	5.9 ± 0.4	16.8 ± 1.2	4.5	7.2	57.1	0	90	42
Augusta	A5560VT3	5.9 ± 0.4	16.8 ± 1.2	5.2	6.5	55.5	0	82	39
Mycogen	TMF2H918 (RR/LL/HX)	5.9 ± 0.4	16.7 ± 1.2	4.2	7.5	67.0	0	94	46
Augusta	A6166GT3000A (RR/LL/CB/RW)	5.8 ± 0.4	16.7 ± 1.2	4.8	6.9	57.9	0	84	41
Augusta	A5337EVT3	5.8 ± 0.4	16.6 ± 1.2	5.4	6.2	61.8	2	87	40
Croplan	9009 RH	5.4 ± 0.4	15.5 ± 1.2	4.3	6.6	65.1	0	90	50
Wyffels	W8437 (VT3Pro)	5.4 ± 0.4	15.3 ± 1.2	4.7	6.0	55.5	0	73	38
Croplan	8221VT3	4.9 ± 0.4	14.0 ± 1.2	3.6	6.2	68.3	0	90	50
Augusta	A5464GTCBLL	4.8 ± 0.4	13.8 ± 1.2	3.8	5.9	61.3	0	84	42
Avg. (tons/a)		5.9	16.8	4.9	6.8	59.4	0	86	44
L.S.D._{.05} (tons/a)		1.3	3.7						
C.V. (%)		10.5	10.5						

† all silage yields are adjusted to dry weight basis unless otherwise indicated.

YG, YGCB, CB, Bt, HX = contains a *Bacillus thuringiensis* gene for insect resistance

YGRW, CRW, RW = contains a gene for rootworm resistance

RR, R, RR2, R2, GT = contains a gene for tolerance to glyphosate

LL = contains a gene for tolerance to glufosinate

CL = contains a gene for tolerance to Imidazolinone class herbicides

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

Blount County Mac Pate Dairy Farm

Planted: 4-25-11

Harvested: 8-1-11

Population: 27,000

30 inch row spacing

Washington County: Savland farm (David Saylor)

Planted: 6-3-11

Harvested: 9-1-11

Population: 28,000

30 inch row spacing

Table 13. Mean yields † and feed quality characteristics of 20 corn hybrids evaluated for silage in two County Standard Tests in Tennessee during 2011.

Brand	Hybrid §	Dry Weight											
		Avg. Yield ± Std Err. (n=2)	Moisture at Harvest (n=2)	Crude Protein (n=2)	NDF 30h			Starch (n=2)	ADF (n=2)	TDN (n=2)	NEL (n=2)	Milk/ton‡ (n=2)	Milk/acre‡ (n=2)
		tons/a	%	% dm	% dm	% of NDF	% dm	% dm	% dm	Mcals/lb	lbs/ton	lbs/acre	
Augusta	A7664VT3	6.7 ± 0.4	50.5	9.8	44.7	64.5	26.5	26.0	71.4	0.74	2928	19556	
DeKalb	DKC67-21 GENVT3P	6.4 ± 0.4	52.3	9.0	52.6	61.2	17.9	31.4	67.3	0.70	2683	17251	
Croplan	7131VT3	6.3 ± 0.4	59.3	9.3	42.9	61.7	29.4	25.0	72.1	0.75	3057	19383	
Augusta	A5462GT3000A (RR/LLCB	6.3 ± 0.4	56.5	8.3	43.4	62.4	30.2	25.3	72.2	0.75	2927	18355	
Croplan	8756VT3	6.1 ± 0.4	63.1	8.8	48.9	62.7	20.9	28.4	69.9	0.73	2957	18186	
Wyffels	W8681 (VT3Pro)	6.0 ± 0.4	56.5	9.5	48.2	60.8	22.8	28.8	69.3	0.72	2797	16921	
Croplan	8505VT3P	6.0 ± 0.4	57.3	8.8	44.2	59.0	28.3	26.3	71.2	0.74	2858	17061	
Croplan	6286 SS (RR2/LL/CB/RW)	6.0 ± 0.4	60.0	8.3	43.7	58.1	30.7	25.3	72.2	0.75	3003	17896	
Masters Choice	MCT-6583 (RR/LL/CB/RW)	5.9 ± 0.4	59.1	8.6	44.0	60.8	29.7	25.6	71.7	0.75	2985	17612	
Augusta	A5461GTCBLLA	5.9 ± 0.4	58.1	8.6	45.5	58.5	27.3	27.5	70.2	0.73	2820	16638	
Mycogen	TMF2L872 (RR/LL/HX1)	5.9 ± 0.4	65.1	8.4	47.4	58.7	20.9	28.7	70.0	0.73	2821	16641	
Augusta	A6867GTCBLLA	5.9 ± 0.4	57.1	9.8	43.5	60.6	28.6	25.7	71.2	0.74	2881	16967	
Augusta	A5560VT3	5.9 ± 0.4	55.5	8.8	45.7	57.7	27.2	27.1	70.7	0.74	2758	16191	
Mycogen	TMF2H918 (RR/LL/HX)	5.9 ± 0.4	67.0	9.3	49.1	59.2	21.7	29.5	69.2	0.72	3113	18210	
Augusta	A6166GT3000A (RR/LL/CL	5.8 ± 0.4	57.9	8.5	46.1	62.3	27.0	26.8	70.9	0.74	2933	17129	
Augusta	A5337EVT3	5.8 ± 0.4	61.8	8.7	43.7	59.1	27.6	26.4	71.1	0.74	3078	17885	
Croplan	9009 RH	5.4 ± 0.4	65.1	9.4	46.3	62.0	24.0	27.7	70.4	0.73	3180	17202	
Wyffels	W8437 (VT3Pro)	5.4 ± 0.4	55.5	9.0	46.7	58.9	25.0	28.0	69.8	0.73	2740	14656	
Croplan	8221VT3	4.9 ± 0.4	68.3	9.3	45.7	57.6	23.1	27.9	70.0	0.73	3045	14889	
Augusta	A5464GTCBLL	4.8 ± 0.4	61.3	9.5	44.1	61.6	29.3	25.7	71.5	0.75	3081	14912	

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

Bt, YG, YGCB, CB, HX = contains a *Bacillus thuringiensis* gene for insect resistance

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

YGRW, RW = contains a gene for rootworm resistance

VT3, TS = contains genes for corn borer, rootworm, and glyphosate resistance

NDF = Neutral Detergent Fiber

VT3P = contains genes for corn borer, rootworm, earworm, armyworm and glyphosate resistance

ADF = Acid Detergent Fiber

RR, RR2 = contains a gene for tolerance to glyphosate

TDN = Total Digestable Nutrients

LL = contains a gene for tolerance to glufosinate

NEL = Net Energy for Lactation

CL = contains a gene for tolerance to Imidazolinone class herbicides

‡ based on University of Wisconsin Milk2006 software program.

Small Grains Silage Test

Seventy-one wheat varieties and 23 fall seeded oat varieties were evaluated for silage yield and quality at the Middle Tennessee REC (Spring Hill, TN). Varieties were seeded at 26 seed per square foot. Plots were planted on 10/19/10 with a drill and consisted of 7 rows on 7 inch spacings, 30 ft in length. Each entry was replicated three times. The plots were harvested on 4/25/11 by a commercial silage harvester. Yields presented were adjusted to both dry weight and 65% moisture. A sub-sample from each plot of approximately 3 lbs was taken for analysis. Fresh weight and dried weight was recorded on each sample for determination of moisture at harvest. The samples were then ground and analyzed for nutrient content. Silage quality analyses were provided by Cumberland Valley Analytical Services, Inc., Hagerstown, MD. Milk per ton and milk per acre calculations were performed using the University of Wisconsin Milk2006 program.

Table 14. Mean yields and agronomic characteristics of 71 soft red winter wheat varieties evaluated for silage at the Middle Tennessee Research and Education Center during 2011.

Brand	Variety	Dry Weight	65% Moisture		Moisture at harvest (n=1)	Emergence (n=1)	Lodging (n=1)	Height (n=1)
		Avg. Yield ± Std Err. (n=1)	Avg. Yield ± Std Err. (n=1)	%				
		tons/a	tons/a	%	%	(score)	inches	
Dyna-Gro	9922	4.8 ± 0.3	13.8 ± 0.9	69.3	100	1.0	33	
Pioneer	XW09H	4.5 ± 0.3	12.7 ± 0.9	68.9	100	1.0	31	
USG	3251	4.4 ± 0.3	12.7 ± 0.9	71.0	97	1.0	33	
Armor	ARX 0186	4.4 ± 0.3	12.5 ± 0.9	69.4	92	1.0	33	
Warren Seed	McKenna 200	4.3 ± 0.3	12.3 ± 0.9	69.0	98	1.0	32	
Pioneer	26R15	4.3 ± 0.3	12.3 ± 0.9	69.0	100	1.0	33	
Syngenta	SY 9978	4.3 ± 0.3	12.2 ± 0.9	69.6	97	1.0	35	
Terral	LA821	4.3 ± 0.3	12.2 ± 0.9	69.7	98	1.0	37	
MO	Truman	4.2 ± 0.3	12.1 ± 0.9	73.2	88	1.0	31	
Progeny	PGX10-7	4.2 ± 0.3	12.1 ± 0.9	70.2	98	1.0	33	
Progeny	PGX10-5	4.2 ± 0.3	12.0 ± 0.9	68.8	100	1.0	33	
Terral	TV8861	4.2 ± 0.3	11.9 ± 0.9	69.7	98	1.0	30	
USG	3438	4.2 ± 0.3	11.9 ± 0.9	69.9	93	1.0	33	
Progeny	125	4.2 ± 0.3	11.9 ± 0.9	69.1	98	1.0	34	
Terral	TVX8535	4.1 ± 0.3	11.7 ± 0.9	72.0	95	1.0	33	
USG	3209	4.1 ± 0.3	11.6 ± 0.9	70.6	97	1.0	32	
TN Exp.	TN 902	4.1 ± 0.3	11.6 ± 0.9	68.8	100	1.0	36	
Pioneer	26R22	4.0 ± 0.3	11.5 ± 0.9	71.5	100	1.0	32	
Pioneer	26R20	4.0 ± 0.3	11.5 ± 0.9	71.6	98	1.0	32	
Terral	TV8589	4.0 ± 0.3	11.4 ± 0.9	70.9	98	1.0	34	
Delta Grow	8300	4.0 ± 0.3	11.3 ± 0.9	69.1	97	1.0	36	
Armor	Ricochet	4.0 ± 0.3	11.3 ± 0.9	72.2	100	1.0	32	
Croplan Genetics	8302	4.0 ± 0.3	11.3 ± 0.9	71.1	97	1.0	34	
USG	3770	3.9 ± 0.3	11.2 ± 0.9	69.3	98	1.0	31	
Armor	ARX 0179	3.9 ± 0.3	11.2 ± 0.9	69.9	97	1.0	35	
MO	Milton	3.9 ± 0.3	11.2 ± 0.9	72.0	88	1.0	34	
Terral	TVX8525	3.9 ± 0.3	11.1 ± 0.9	70.8	97	1.0	33	
Delta Grow	7900	3.9 ± 0.3	11.0 ± 0.9	69.5	100	1.0	35	
OH	Malabar	3.9 ± 0.3	11.0 ± 0.9	73.2	100	1.0	30	
Cache River Valley Seed	Dixie McAlister	3.8 ± 0.3	11.0 ± 0.9	68.6	97	1.0	33	
USG	3120	3.8 ± 0.3	10.9 ± 0.9	69.9	97	1.0	35	
USG	3555	3.8 ± 0.3	10.8 ± 0.9	71.0	100	1.0	32	
VA Exp.	VA05W-251	3.8 ± 0.3	10.7 ± 0.9	70.4	97	1.0	33	
Progeny	166	3.8 ± 0.3	10.7 ± 0.9	69.4	100	1.0	34	
Delta Grow	7500	3.8 ± 0.3	10.7 ± 0.9	69.8	97	1.0	34	
Progeny	PGX10-2	3.7 ± 0.3	10.6 ± 0.9	71.1	98	1.0	33	
TN Exp.	TN 1002	3.7 ± 0.3	10.6 ± 0.9	70.3	97	1.0	37	
Warren Seed	McKay 100	3.7 ± 0.3	10.6 ± 0.9	72.4	100	1.0	35	
Armor	Renegade	3.7 ± 0.3	10.6 ± 0.9	71.8	95	1.0	32	
NC Exp.	Yadkin	3.7 ± 0.3	10.5 ± 0.9	69.2	93	1.0	32	
Croplan Genetics	8614	3.7 ± 0.3	10.5 ± 0.9	69.0	97	1.0	36	
Terral	TVX8626	3.7 ± 0.3	10.4 ± 0.9	71.9	100	1.0	32	
USG	3201	3.7 ± 0.3	10.4 ± 0.9	70.9	97	1.0	31	
Dyna-Gro	9171	3.6 ± 0.3	10.4 ± 0.9	69.1	92	1.0	32	
USG	3295	3.6 ± 0.3	10.4 ± 0.9	68.1	98	1.0	31	
Progeny	185	3.6 ± 0.3	10.3 ± 0.9	71.2	100	1.0	33	
Terral	TV8558	3.6 ± 0.3	10.2 ± 0.9	70.8	93	1.0	33	
Syngenta	Oakes	3.6 ± 0.3	10.2 ± 0.9	70.9	100	1.0	31	
Syngenta	W1104	3.6 ± 0.3	10.2 ± 0.9	70.3	92	1.0	30	
Delta Grow	5000	3.6 ± 0.3	10.1 ± 0.9	71.1	98	1.0	35	
VA Exp.	VA05W-139	3.6 ± 0.3	10.1 ± 0.9	70.9	92	1.0	31	

(continued)

Table 14. Mean yields and agronomic characteristics of 71 soft red winter wheat varieties evaluated for silage at the Middle Tennessee Research and Education Center during 2011.

Brand	Variety	Dry Weight		65% Moisture		Emergence (n=1)	Lodging (n=1)	Height (n=1)
		Avg. Yield ± Std Err. (n=1)	Avg. Yield ± Std Err. (n=1)	Moisture at harvest (n=1)	%			
TN Exp.	TN 1103	3.5 ± 0.3	10.1 ± 0.9	70.6	100	1.0	33	
USG	3409	3.5 ± 0.3	10.0 ± 0.9	70.1	98	1.0	34	
Cache River Valley Seed	Dixie Kelsey	3.5 ± 0.3	10.0 ± 0.9	69.8	98	1.0	32	
Dyna-Gro	9012	3.5 ± 0.3	10.0 ± 0.9	71.5	97	1.0	31	
MO	Bess	3.4 ± 0.3	9.8 ± 0.9	70.8	98	1.0	32	
VA	Jamestown	3.4 ± 0.3	9.8 ± 0.9	71.2	100	1.0	35	
Cache River Valley Seed	Dixie 454	3.4 ± 0.3	9.7 ± 0.9	70.4	97	1.0	31	
Syngenta	Branson	3.4 ± 0.3	9.7 ± 0.9	69.2	95	1.0	33	
Pioneer	25R32	3.4 ± 0.3	9.6 ± 0.9	70.5	97	1.0	30	
VA	Merl	3.3 ± 0.3	9.4 ± 0.9	72.3	100	1.0	34	
TN Exp.	TN 1101	3.3 ± 0.3	9.3 ± 0.9	71.3	98	1.0	37	
USG	3350	3.2 ± 0.4	9.3 ± 1.1	71.4	95	1.0	34	
Terral	TVX8848	3.2 ± 0.3	9.3 ± 0.9	71.9	68	1.0	32	
Dyna-Gro	Shirley	3.2 ± 0.3	9.2 ± 0.9	71.1	97	1.0	31	
Progeny	117	3.2 ± 0.3	9.2 ± 0.9	75.5	100	1.0	36	
Croplan Genetics	8925	3.2 ± 0.3	9.1 ± 0.9	71.3	97	1.0	34	
TN Exp.	TN 1102	3.2 ± 0.3	9.0 ± 0.9	73.4	100	1.0	35	
USG	3345	3.1 ± 0.3	8.9 ± 0.9	70.6	92	1.0	33	
Terral	TVX8460	3.1 ± 0.3	8.8 ± 0.9	73.9	72	1.0	30	
Dyna-Gro	9053	2.6 ± 0.3	7.5 ± 0.9	76.2	98	1.0	32	
		Average (bu/a)	3.8	10.7	70.7	96	1.0	33
		L.S.D._{.05} (bu/a)	0.8	2.3				
		C.V. (%)	13.2	13.2				

Lodging = 1 to 5 scale; where 1 = 95% of plants erect; 2.5 = ~50% of plants leaning at angle $\geq 45^\circ$; 5 = 95+% of plants leaning at an angle $\geq 45^\circ$.

Table 15. Mean yields † and feed quality characteristics of 71 wheat varieties evaluated for silage at the Middle Tennessee Research and Education Center during 2011.

Brand	Variety	Dry Weight											
		Avg. Yield ± Std Err. (n=1)	Moisture at Harvest (n=1)	Crude Protein (n=1)	NDF 30h								
		%	% dm	% dm	NDF (n=1)	IV Digest (n=1)	Starch (n=1)	ADF (n=1)	TDN (n=1)	NEL (n=1)	Milk/ton‡ (n=1)	Milk/acre‡ (n=1)	
Dyna-Gro	9922	tons/a 4.8 ± 0.3	% 69.3	% dm 9.1	% dm 62.2	% of NDF 60.1	% dm 3.1	% dm 39.6	% dm 59.6	Mcals/lb 0.61	lbs/ton 2439	lbs/acre 11732	
Pioneer	XW09H	4.5 ± 0.3	68.9	8.8	62.2	56.8	2.9	39.7	59.5	0.61	2255	10035	
USG	3251	4.4 ± 0.3	71.0	9.3	65.7	60.4	1.5	41.5	58.3	0.60	2509	11115	
Armor	ARX 0186	4.4 ± 0.3	69.4	9.3	58.3	58.8	3.6	36.8	61.7	0.64	2270	9896	
Warren Seed	McKenna 200	4.3 ± 0.3	69.0	9.2	65.5	56.9	1.1	42.4	56.0	0.57	2281	9831	
Pioneer	26R15	4.3 ± 0.3	69.0	8.8	61.3	58.2	3.1	39.1	59.6	0.61	2304	9886	
Syngenta	SY 9978	4.3 ± 0.3	69.6	7.7	62.9	56.9	2.1	40.9	57.6	0.59	2118	9064	
Terral	LA821	4.3 ± 0.3	69.7	8.2	66.2	56.2	1.7	42.3	57.3	0.59	2260	9629	
MO	Truman	4.2 ± 0.3	73.2	11.9	63.9	62.8	2.9	39.4	60.6	0.62	2697	11437	
Progeny	PGX10-7	4.2 ± 0.3	70.2	9.9	68.7	55.4	2.4	44.2	56.5	0.58	2276	9606	
Progeny	PGX10-5	4.2 ± 0.3	68.8	8.7	61.9	56.4	2.3	39.8	58.1	0.60	2181	9180	
Terral	TV8861	4.2 ± 0.3	69.7	8.2	60.9	60.6	2.3	38.6	60.0	0.62	2228	9311	
USG	3438	4.2 ± 0.3	69.9	7.8	59.0	59.0	3.1	37.6	59.9	0.61	2105	8799	
Progeny	125	4.2 ± 0.3	69.1	
Terral	TVX8535	4.1 ± 0.3	72.0	9.3	68.8	53.1	2.3	44.6	55.1	0.56	2124	8664	
USG	3209	4.1 ± 0.3	70.6	8.7	67.4	56.2	1.5	43.2	56.6	0.58	2351	9567	
TN Exp.	TN 902	4.1 ± 0.3	68.8	9.2	59.6	59.5	2.6	38.0	59.5	0.61	2259	9150	
Pioneer	26R22	4.0 ± 0.3	71.5	9.2	61.4	62.6	2.5	39.2	61.0	0.63	2441	9836	
Pioneer	26R20	4.0 ± 0.3	71.6	8.2	61.8	61.3	2.8	39.5	59.1	0.61	2364	9481	
Terral	TV8589	4.0 ± 0.3	70.9	7.9	61.7	58.0	2.1	39.5	58.5	0.60	2151	8602	
Delta Grow	8300	4.0 ± 0.3	69.1	7.8	62.9	58.0	3.5	39.7	59.0	0.61	2310	9169	
Armor	Ricochet	4.0 ± 0.3	72.2	8.7	71.2	48.5	3.9	46.6	53.6	0.54	1831	7252	
Croplan Genetics	8302	4.0 ± 0.3	71.1	8.3	61.1	58.3	3.1	38.8	59.5	0.61	2239	8844	
USG	3770	3.9 ± 0.3	69.3	7.9	61.5	55.4	2.8	39.2	58.7	0.60	2059	8093	
Armor	ARX 0179	3.9 ± 0.3	69.9	8.8	71.5	49.2	3.8	46.6	54.5	0.55	1937	7574	
MO	Milton	3.9 ± 0.3	72.0	7.9	66.8	55.7	2.7	43.0	57.5	0.59	2353	9175	
Terral	TVX8525	3.9 ± 0.3	70.8	8.8	59.2	62.4	2.5	37.2	60.8	0.62	2272	8794	
Delta Grow	7900	3.9 ± 0.3	69.5	8.9	71.8	48.1	3.9	47.0	53.7	0.54	1839	7099	
OH	Malabar	3.9 ± 0.3	73.2	10.2	61.1	66.4	2.1	37.5	61.2	0.63	2621	10117	
Cache River Valley Seed	Dixie McAlister	3.8 ± 0.3	68.6	8.6	64.7	58.5	2.9	41.1	58.4	0.60	2446	9391	
USG	3120	3.8 ± 0.3	69.9	9.8	59.8	60.7	1.8	38.1	59.1	0.61	2280	8731	
USG	3555	3.8 ± 0.3	71.0	8.1	65.6	61.8	2.3	40.6	59.2	0.61	2479	9370	
VA Exp.	VA05W-251	3.8 ± 0.3	70.4	8.7	66.8	57.3	1.9	43.0	56.4	0.58	2380	8950	
Progeny	166	3.8 ± 0.3	69.4	8.4	69.0	56.1	1.7	44.6	55.8	0.57	2288	8579	
Delta Grow	7500	3.8 ± 0.3	69.8	8.8	57.5	59.4	3.3	36.3	61.5	0.63	2160	8098	
Progeny	PGX10-2	3.7 ± 0.3	71.1	9.1	64.8	59.6	1.8	42.0	57.7	0.59	2431	9042	

Table 15. (continued)

Brand	Variety	Dry Weight													
		Avg. Yield		Moisture at Harvest	Crude		NDF 30h						NEL (n=1)	Milk/ton [‡] (n=1)	Milk/acre [‡] (n=1)
		± Std Err. (n=1)	(n=1)		Protein (n=1)	NDF (n=1)	IV Digest (n=1)	Starch (n=1)	ADF (n=1)	TDN (n=1)	Mcals/lb	lbs/ton			
		tons/a	%	% dm	% dm	% of NDF	% dm	% dm	% dm	% dm					
TN Exp.	TN 1002	3.7 ± 0.3	70.3	8.3	65.2	58.6	2.2	41.6	57.5	0.59	2355	8760			
Warren Seed	McKay 100	3.7 ± 0.3	72.4	8.6	64.5	56.7	1.7	41.6	56.8	0.58	2240	8311			
Armor	Renegade	3.7 ± 0.3	71.8	8.4	60.8	55.6	3.4	39.2	59.7	0.61	2196	8124			
NC Exp.	Yadkin	3.7 ± 0.3	69.2	7.8	63.1	57.7	2.8	40.6	58.8	0.60	2237	8212			
Croplan Genetics	8614	3.7 ± 0.3	69.0	8.5	62.2	59.3	3.2	39.3	60.1	0.62	2332	8536			
Terral	TVX8626	3.7 ± 0.3	71.9	9.1	64.2	60.0	2.1	40.4	59.0	0.61	2424	8849			
USG	3201	3.7 ± 0.3	70.9	9.3	64.3	60.8	2.1	40.5	58.1	0.59	2440	8907			
Dyna-Gro	9171	3.6 ± 0.3	69.1	9.2	67.1	54.7	3.3	42.2	57.2	0.59	2303	8383			
USG	3295	3.6 ± 0.3	68.1	8.7	62.9	59.7	1.4	39.9	58.6	0.60	2255	8207			
Progeny	185	3.6 ± 0.3	71.2	10.1	65.1	60.2	1.5	41.7	58.1	0.59	2530	9084			
Terral	TV8558	3.6 ± 0.3	70.8	9.3	68.6	53.1	2.8	43.8	55.5	0.57	2179	7778			
Syngenta	Oakes	3.6 ± 0.3	70.9	10.0	63.9	59.0	1.3	40.6	58.2	0.60	2437	8677			
Syngenta	W1104	3.6 ± 0.3	70.3	8.6	74.1	51.7	1.4	47.9	52.9	0.54	1944	6922			
Delta Grow	5000	3.6 ± 0.3	71.1	9.3	71.8	49.6	5.0	44.6	55.0	0.56	1997	7091			
VA Exp.	VA05W-139	3.6 ± 0.3	70.9	8.7	63.6	60.8	2.2	40.0	59.6	0.61	2398	8513			
TN Exp.	TN 1103	3.5 ± 0.3	70.6	9.5	61.9	57.4	2.1	39.6	57.9	0.59	2250	7943			
USG	3409	3.5 ± 0.3	70.1	8.9	62.1	54.7	2.2	39.3	58.7	0.60	2132	7485			
Cache River Valley Seed	Dixie Kelsey	3.5 ± 0.3	69.8	8.9	62.7	59.9	3.0	39.4	59.8	0.61	2388	8383			
Dyna-Gro	9012	3.5 ± 0.3	71.5	8.8	59.6	60.2	3.7	37.7	60.8	0.62	2292	7998			
MO	Bess	3.4 ± 0.3	70.8	9.4	63.0	59.1	3.4	40.0	59.4	0.61	2514	8649			
VA	Jamestown	3.4 ± 0.3	71.2	9.4	62.4	56.9	2.5	39.5	58.9	0.60	2283	7832			
Cache River Valley Seed	Dixie 454	3.4 ± 0.3	70.4	8.3	57.9	59.5	4.1	36.7	61.2	0.63	2198	7495			
Syngenta	Branson	3.4 ± 0.3	69.2	8.7	61.3	58.1	1.5	39.6	57.7	0.59	2131	7224			
Pioneer	25R32	3.4 ± 0.3	70.5	9.0	69.0	58.9	2.5	43.8	57.0	0.58	2409	8069			
VA	Merl	3.3 ± 0.3	72.3	8.5	70.3	49.8	2.7	45.3	54.6	0.56	2024	6678			
TN Exp.	TN 1101	3.3 ± 0.3	71.3	9.1	71.4	54.6	1.4	46.2	52.8	0.54	2019	6582			
USG	3350	3.2 ± 0.4	71.4	8.7	67.6	56.5	1.9	43.2	56.6	0.58	2381	7714			
Terral	TVX8848	3.2 ± 0.3	71.9	8.4	63.3	59.6	1.9	40.2	58.6	0.60	2304	7465			
Dyna-Gro	Shirley	3.2 ± 0.3	71.1	8.3	63.4	57.1	3.3	40.6	58.5	0.60	2316	7457			
Progeny	117	3.2 ± 0.3	75.5	7.9	63.6	57.7	1.8	41.2	56.9	0.58	2186	7019			
Croplan Genetics	8925	3.2 ± 0.3	71.3	8.9	62.6	61.6	3.2	39.2	61.0	0.63	2478	7905			
TN Exp.	TN 1102	3.2 ± 0.3	73.4	9.4	66.9	58.7	1.8	42.5	56.7	0.58	2409	7588			
USG	3345	3.1 ± 0.3	70.6	8.9	65.6	56.9	2.1	41.6	57.7	0.59	2323	7248			
Terral	TVX8460	3.1 ± 0.3	73.9	9.3	59.1	63.4	2.6	37.2	60.8	0.63	2402	7421			
Dyna-Gro	9053	2.6 ± 0.3	76.2	9.0	66.1	53.7	3.4	42.6	57.0	0.58	2254	5884			

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

Table 16. Mean yields and agronomic characteristics of 44 soft red winter wheat varieties evaluated for silage at the Middle Tennessee Research and Education Center for two years (2010-2011).

Brand	Variety	Dry Weight	65% Moisture				Lodging (n=2)	Height (n=2)
		Avg. Yield ± Std Err. (n=2)	Avg. Yield ± Std Err. (n=2)	Moisture at harvest (n=2)	Emergence (n=1)			
USG	3251	4.0 ± 0.2	11.4 ± 0.6	65.3	97	1.0	32	
Croplan Genetics	8302	3.9 ± 0.2	11.3 ± 0.6	63.0	97	1.0	33	
Syngenta	SY 9978	3.9 ± 0.2	11.1 ± 0.6	65.5	97	1.0	36	
TN Exp.	TN 902	3.8 ± 0.2	10.9 ± 0.6	63.2	100	1.0	34	
Dyna-Gro	9922	3.8 ± 0.2	10.8 ± 0.6	63.1	100	1.0	32	
Pioneer	26R22	3.8 ± 0.2	10.8 ± 0.6	64.4	100	1.0	32	
Terral	LA821	3.7 ± 0.2	10.7 ± 0.6	62.3	98	1.0	35	
Armor	Ricochet	3.7 ± 0.2	10.6 ± 0.6	64.1	100	1.0	30	
Progeny	125	3.7 ± 0.2	10.6 ± 0.6	61.6	98	1.0	31	
Pioneer	26R15	3.7 ± 0.2	10.6 ± 0.6	65.6	100	1.0	32	
USG	3209	3.7 ± 0.2	10.4 ± 0.6	64.1	97	1.0	30	
USG	3438	3.6 ± 0.2	10.4 ± 0.6	65.2	93	1.0	30	
USG	3770	3.6 ± 0.2	10.3 ± 0.6	62.9	98	1.0	31	
Delta Grow	8300	3.6 ± 0.2	10.1 ± 0.6	63.4	97	1.0	33	
Syngenta	W1104	3.5 ± 0.2	10.1 ± 0.6	65.2	92	1.0	31	
MO	Truman	3.5 ± 0.2	10.1 ± 0.6	67.3	88	1.0	32	
Terral	TV8589	3.5 ± 0.2	10.0 ± 0.6	64.5	98	1.0	33	
Warren Seed	McKay 100	3.5 ± 0.2	10.0 ± 0.6	63.2	100	1.0	34	
USG	3201	3.5 ± 0.2	9.9 ± 0.6	63.5	97	1.0	30	
Pioneer	26R20	3.5 ± 0.2	9.9 ± 0.6	68.3	98	1.0	32	
MO	Milton	3.4 ± 0.2	9.8 ± 0.6	64.9	88	1.0	32	
Progeny	166	3.4 ± 0.2	9.8 ± 0.6	63.9	100	1.0	34	
Dyna-Gro	9012	3.4 ± 0.2	9.7 ± 0.6	65.6	97	1.0	30	
Terral	TV8558	3.4 ± 0.2	9.7 ± 0.6	62.3	93	1.0	32	
USG	3120	3.4 ± 0.2	9.6 ± 0.6	63.8	97	1.0	34	
OH	Malabar	3.3 ± 0.2	9.6 ± 0.6	68.4	100	1.0	31	
USG	3350	3.3 ± 0.2	9.5 ± 0.6	64.8	95	1.0	34	
USG	3555	3.3 ± 0.2	9.5 ± 0.6	65.9	100	1.0	28	
MO	Bess	3.3 ± 0.2	9.4 ± 0.6	63.6	98	1.0	32	
Armor	Renegade	3.3 ± 0.2	9.4 ± 0.6	64.5	95	1.0	32	
Terral	TV8861	3.3 ± 0.2	9.4 ± 0.6	64.4	98	1.0	30	
USG	3409	3.2 ± 0.2	9.3 ± 0.6	63.5	98	1.0	32	
Delta Grow	5000	3.2 ± 0.2	9.2 ± 0.6	63.8	98	1.0	32	
Progeny	185	3.2 ± 0.2	9.1 ± 0.6	64.6	100	1.0	31	
Syngenta	Oakes	3.2 ± 0.2	9.1 ± 0.6	65.5	100	1.0	31	
Progeny	117	3.1 ± 0.2	9.0 ± 0.6	67.1	100	1.0	34	
VA	Merl	3.1 ± 0.2	8.9 ± 0.6	64.9	100	1.0	31	
Cache River Valley Seed	Dixie 454	3.1 ± 0.2	8.9 ± 0.6	66.0	97	1.0	31	
Croplan Genetics	8925	3.1 ± 0.2	8.9 ± 0.6	63.7	97	1.0	33	
Dyna-Gro	Shirley	3.1 ± 0.2	8.8 ± 0.6	65.4	97	1.0	28	
NC Exp.	Yadkin	3.1 ± 0.2	8.8 ± 0.6	64.2	93	1.0	30	
VA	Jamestown	3.1 ± 0.2	8.7 ± 0.6	64.4	100	1.0	31	
Pioneer	25R32	3.0 ± 0.2	8.5 ± 0.6	66.0	97	1.0	31	
Syngenta	Branson	2.9 ± 0.2	8.4 ± 0.6	65.2	95	1.0	31	
		Average (bu/a)	3.4	9.8	64.6	97	1.0	32
		L.S.D._{.05} (bu/a)	0.6	1.8				
		C.V. (%)	12.9	12.9				

Lodging = 1 to 5 scale; where 1 = 95% of plants erect; 2.5 = ~50% of plants leaning at angle ≥ 45°; 5 = 95+% of plants leaning at an angle ≥ 45°.

Table 17. Mean yields † and feed quality characteristics of 44 wheat varieties evaluated for silage at the Middle Tennessee Research and Education Center for two years (2010-2011).

Brand	Variety	Dry Weight											
		Avg. Yield ± Std Err. (n=2)	Moisture at Harvest (n=2)	Crude Protein (n=2)	NDF (n=2)	NDF 30h		Starch (n=2)	ADF (n=2)	TDN (n=2)	NEL (n=2)	Milk/ton‡ (n=2)	Milk/acre‡ (n=2)
		%	% dm	% dm	% of NDF	IV Digest (n=2)	% dm					Mcals/lb	lbs/ton
USG	3251	4.0 ± 0.2	65.3	8.8	59.5	56.9	5.5	37.3	57.4	0.56	2323	9392	
Croplan Genetics	8302	3.9 ± 0.2	63.0	8.4	58.6	56.0	4.6	36.9	57.2	0.56	2120	8353	
Syngenta	SY 9978	3.9 ± 0.2	65.5	8.7	59.6	54.5	5.0	38.4	57.8	0.57	2196	8467	
TN Exp.	TN 902	3.8 ± 0.2	63.2	8.0	57.7	57.5	4.9	36.4	57.6	0.56	2146	8225	
Dyna-Gro	9922	3.8 ± 0.2	63.1	8.7	57.2	56.4	7.3	36.3	58.2	0.57	2298	8778	
Pioneer	26R22	3.8 ± 0.2	64.4	9.4	58.9	58.7	5.5	37.5	60.0	0.59	2374	8955	
Terral	LA821	3.7 ± 0.2	62.3	7.5	62.5	53.5	4.4	39.5	55.3	0.54	2088	7888	
Armor	Ricochet	3.7 ± 0.2	64.1	9.1	62.9	51.7	5.6	40.3	55.1	0.53	1971	7319	
Progeny	125	3.7 ± 0.2	61.6	9.0	56.6	52.2	7.2	35.3	55.6	0.52	2079	6818	
Pioneer	26R15	3.7 ± 0.2	65.6	9.3	59.7	55.5	5.8	37.8	59.5	0.59	2344	8613	
USG	3209	3.7 ± 0.2	64.1	8.3	60.2	57.0	4.9	38.0	57.0	0.55	2245	8217	
USG	3438	3.6 ± 0.2	65.2	8.8	56.3	57.2	6.3	35.4	59.8	0.59	2230	7990	
USG	3770	3.6 ± 0.2	62.9	8.2	58.3	55.4	5.8	36.8	58.4	0.57	2146	7675	
Delta Grow	8300	3.6 ± 0.2	63.4	8.3	58.3	56.4	6.8	36.7	58.8	0.58	2292	8146	
Syngenta	W1104	3.5 ± 0.2	65.2	9.4	65.5	53.1	3.8	41.9	55.8	0.55	2115	7437	
MO	Truman	3.5 ± 0.2	67.3	11.0	61.3	59.5	3.3	38.0	59.4	0.58	2463	8840	
Terral	TV8589	3.5 ± 0.2	64.5	8.1	59.9	54.6	5.0	37.7	57.4	0.57	2150	7557	
Warren Seed	McKay 100	3.5 ± 0.2	63.2	8.6	60.3	56.2	4.9	38.2	57.3	0.56	2222	7759	
USG	3201	3.5 ± 0.2	63.5	8.4	58.7	58.5	4.8	36.8	56.3	0.54	2187	7634	
Pioneer	26R20	3.5 ± 0.2	68.3	8.8	59.6	60.1	5.3	37.4	60.7	0.60	2440	8412	
MO	Milton	3.4 ± 0.2	64.9	8.3	61.0	56.1	4.6	38.6	57.1	0.56	2227	7676	
Progeny	166	3.4 ± 0.2	63.9	8.8	62.4	55.4	5.2	39.7	57.2	0.56	2287	7832	
Dyna-Gro	9012	3.4 ± 0.2	65.6	8.7	57.0	58.1	5.9	35.9	59.2	0.58	2237	7611	
Terral	TV8558	3.4 ± 0.2	62.3	8.1	63.6	52.4	5.0	40.3	54.7	0.54	2067	7036	
USG	3120	3.4 ± 0.2	63.8	8.6	57.2	58.0	5.2	36.1	57.8	0.57	2186	7432	
OH	Malabar	3.3 ± 0.2	68.4	10.4	61.3	62.0	2.8	38.2	61.2	0.60	2531	8489	
USG	3350	3.3 ± 0.2	64.8	8.3	62.1	55.6	4.6	39.3	56.6	0.55	2252	7529	
USG	3555	3.3 ± 0.2	65.9	8.2	59.5	59.0	4.7	37.1	57.9	0.56	2284	7672	
MO	Bess	3.3 ± 0.2	63.6	9.1	59.3	57.2	4.9	37.3	58.0	0.57	2309	7586	
Armor	Renegade	3.3 ± 0.2	64.5	9.0	58.7	54.2	6.3	37.5	59.1	0.58	2251	7326	
Terral	TV8861	3.3 ± 0.2	64.4	8.3	58.1	58.0	4.6	36.5	58.3	0.57	2169	7124	
USG	3409	3.2 ± 0.2	63.5	8.7	58.8	53.8	5.8	36.8	57.9	0.57	2163	6967	
Delta Grow	5000	3.2 ± 0.2	63.8	9.0	62.4	50.8	7.7	38.8	55.7	0.55	2067	6664	
Progeny	185	3.2 ± 0.2	64.6	9.3	61.8	55.7	4.1	39.5	56.8	0.56	2309	7497	
Syngenta	Oakes	3.2 ± 0.2	65.5	9.9	59.8	55.6	4.3	37.8	57.6	0.57	2312	7421	
Progeny	117	3.1 ± 0.2	67.1	8.5	59.6	56.4	4.8	38.2	57.4	0.56	2200	6907	

(continued)

Table 17. Mean yields † and feed quality characteristics of 44 wheat varieties evaluated for silage at the Middle Tennessee Research and Education Center for two years (2010-2011).

Brand	Variety	Dry Weight												
		Avg. Yield ± Std Err. (n=2)	Moisture at Harvest (n=2)	Crude		NDF 30h						NEL (n=2)	Milk/ton‡ (n=2)	Milk/acre‡ (n=2)
				Protein (n=2)	NDF (n=2)	IV Digest (n=2)	Starch (n=2)	ADF (n=2)	TDN (n=2)	% dm	Mcals/lb			
		tons/a	%	% dm	% dm	% of NDF	% dm	% dm	% dm	% dm	lbs/ton	lbs/acre		
VA	Merl	3.1 ± 0.2	64.9	8.3	62.6	52.3	5.4	39.5	55.7	0.54	2075	6486		
Cache River Valley Seed	Dixie 454	3.1 ± 0.2	66.0	9.7	56.0	56.9	6.1	35.2	60.2	0.59	2269	7046		
Croplan Genetics	8925	3.1 ± 0.2	63.7	8.1	59.1	56.7	6.1	37.1	57.9	0.57	2248	7029		
Dyna-Gro	Shirley	3.1 ± 0.2	65.4	8.6	58.2	57.3	4.5	36.9	57.3	0.55	2178	6747		
NC Exp.	Yadkin	3.1 ± 0.2	64.2	8.7	59.6	55.1	4.8	38.3	57.5	0.56	2184	6779		
VA	Jamestown	3.1 ± 0.2	64.4	9.1	57.5	54.8	7.5	35.9	58.5	0.58	2279	6976		
Pioneer	25R32	3.0 ± 0.2	66.0	9.8	63.0	57.1	4.6	39.7	58.2	0.57	2376	7082		
Syngenta	Branson	2.9 ± 0.2	65.2	10.0	57.3	55.6	5.4	36.4	58.7	0.58	2260	6575		

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

Table 18. Mean yields and agronomic characteristics of 23 fall seeded oat lines evaluated for silage at the Middle Tennessee Research and Education Center during 2011.

Brand	Variety	Dry Weight	65% Moisture	Moisture at harvest (n=1)	Emergence (n=1)	Winter Survival (n=1)	Lodging (n=1)	Height (n=1)
		Avg. Yield ± Std Err. (n=1)	Avg. Yield ± Std Err. (n=1)					
		tons/a	tons/a	%	%	%	(score)	inches
TX	TAMO 406	4.6 ± 0.2	13.2 ± 0.6	71.3	93.3	100	1.0	36
TX	TX02D079	4.4 ± 0.2	12.5 ± 0.6	73.4	100.0	100	1.0	35
LA	LA03063SBSBSB-S4	4.4 ± 0.2	12.4 ± 0.6	70.3	96.7	100	1.0	38
LA	LA06041SBS-42	4.3 ± 0.2	12.2 ± 0.6	76.4	98.3	100	1.0	38
TX	TX05CS556	4.0 ± 0.2	11.3 ± 0.6	75.5	100.0	100	1.0	36
LA	LA06059SBS-66	3.9 ± 0.2	11.3 ± 0.6	74.1	96.7	100	1.0	40
NC	NC07-3966	3.9 ± 0.2	11.2 ± 0.6	74.5	96.7	100	1.0	34
LA	Horizon 270	3.9 ± 0.2	11.0 ± 0.6	75.0	98.3	100	1.0	35
LA	LA05006GSBS-65-S1	3.8 ± 0.2	10.9 ± 0.6	77.4	98.3	100	1.0	38
TX	TX09CS1112	3.8 ± 0.2	10.9 ± 0.6	73.3	98.3	100	1.0	32
FL	Horizon 201	3.8 ± 0.2	10.8 ± 0.6	75.2	93.3	100	1.0	40
NC	NC07-3834	3.7 ± 0.2	10.7 ± 0.6	77.2	96.7	100	1.0	35
NC	NC07-3801	3.7 ± 0.2	10.5 ± 0.6	77.2	96.7	100	1.0	37
FL	LA02012-S-B-139-S2-B-S1	3.6 ± 0.2	10.3 ± 0.6	76.0	96.7	100	1.0	39
TX	TX07CS3697	3.6 ± 0.2	10.3 ± 0.6	74.9	100.0	100	1.0	38
FL	FL99913-J1-S1	3.5 ± 0.2	10.1 ± 0.6	76.5	100.0	100	1.0	33
NC	Rodgers	3.4 ± 0.2	9.8 ± 0.6	75.3	86.7	100	1.0	39
TX	TX09CS1025	3.4 ± 0.2	9.8 ± 0.6	75.9	86.7	100	1.0	35
FL	FL04178-FLID-B-S-2	3.4 ± 0.2	9.8 ± 0.6	76.2	98.3	100	1.0	39
TX	TX07CS2783	3.4 ± 0.2	9.7 ± 0.6	76.7	98.3	100	1.0	38
LA	FL0522-FLID-B-S-B-S-92-S1	3.4 ± 0.2	9.6 ± 0.6	77.0	96.7	100	1.0	36
TX	TX09CS1066	3.3 ± 0.2	9.4 ± 0.6	75.3	85.0	100	1.0	34
FL	FL02011-I-J2	2.4 ± 0.2	6.9 ± 0.6	74.8	96.7	62	1.0	35
Average (bu/a)		3.7	10.6	75.2	96.0	98	1.0	37
L.S.D._{.05} (bu/a)		0.6	1.6					
C.V. (%)		9.2	9.2					

Lodging = 1 to 5 scale; where 1 = 95% of plants erect; 2.5 = ~50% of plants leaning at angle $\geq 45^\circ$; 5 = 95+% of plants leaning at an angle $\geq 45^\circ$.

Table 19. Mean yields † and feed quality characteristics of 23 oat varieties evaluated for silage at the Middle Tennessee Research and Education Center during 2011.

Brand	Variety	Dry Weight		Crude Protein (n=1)	NDF 30h						NEL (n=1)	Milk/ton‡ (n=1)	Milk/acre‡ (n=1)
		Avg. Yield ± Std Err. (n=1)	Moisture at Harvest (n=1)		NDF (n=1)	IV Digest (n=1)	Starch (n=1)	ADF (n=1)	TDN (n=1)	Mcals/lb			
		tons/a	%	% dm	% dm	% of NDF	% dm	% dm	% dm	Mcal/lb	lbs/ton	lbs/acre	
TX	TAMO 406	4.6 ± 0.2	71.3	8.3	70.1	58.8	2.2	44.6	56.8	0.58	2190	10075	
TX	TX02D079	4.4 ± 0.2	73.4	10.2	65.2	62.0	0.9	41.4	58.2	0.60	2353	10329	
LA	LA03063SBSBSB-S4	4.4 ± 0.2	70.3	10.9	63.3	62.5	1.6	40.4	59.6	0.61	2462	10712	
LA	LA06041SBS-42	4.3 ± 0.2	76.4	10.4	68.1	57.6	0.9	44.1	55.5	0.57	2047	8740	
TX	TX05CS556	4.0 ± 0.2	75.5	8.4	70.0	51.1	1.9	45.6	56.5	0.58	1979	7856	
LA	LA06059SBS-66	3.9 ± 0.2	74.1	8.2	70.5	60.7	1.2	44.7	57.3	0.59	2308	9092	
NC	NC07-3966	3.9 ± 0.2	74.5	8.9	64.2	58.7	1.8	39.9	60.0	0.62	2237	8745	
LA	Horizon 270	3.9 ± 0.2	75.0	9.1	65.7	58.7	2.0	41.9	57.7	0.59	2286	8823	
LA	LA05006GSBS-65-S1	3.8 ± 0.2	77.4	9.8	62.5	61.3	1.5	39.3	60.0	0.62	2293	8759	
TX	TX09CS1112	3.8 ± 0.2	73.3	9.5	65.6	61.7	1.6	41.9	58.8	0.60	2411	9186	
FL	Horizon 201	3.8 ± 0.2	75.2	7.9	63.9	60.7	2.0	40.9	59.0	0.61	2183	8274	
NC	NC07-3834	3.7 ± 0.2	77.2	9.0	66.9	59.6	1.2	42.6	57.7	0.59	2271	8471	
NC	NC07-3801	3.7 ± 0.2	77.2	9.5	64.4	63.8	1.7	40.2	59.4	0.61	2457	9065	
FL	LA02012-S-B-139-S2-B-S1	3.6 ± 0.2	76.0	9.2	60.6	66.3	1.4	38.6	60.8	0.63	2311	8344	
TX	TX07CS3697	3.6 ± 0.2	74.9	10.1	69.4	55.0	1.9	44.7	55.7	0.57	2043	7375	
FL	FL99913-J1-S1	3.5 ± 0.2	76.5	10.3	63.5	61.2	1.9	40.1	60.2	0.62	2422	8525	
NC	Rodgers	3.4 ± 0.2	75.3	7.9	66.4	64.8	2.3	41.9	59.5	0.61	2475	8463	
TX	TX09CS1025	3.4 ± 0.2	75.9	10.3	66.2	59.2	1.8	41.5	57.5	0.59	2252	7701	
FL	FL04178-FLID-B-S-2	3.4 ± 0.2	76.2	8.0	66.6	56.8	1.4	43.2	57.3	0.59	2150	7330	
TX	TX07CS2783	3.4 ± 0.2	76.7	10.6	61.7	61.7	2.6	38.6	60.3	0.62	2432	8294	
LA	FL0522-FLID-B-S-B-S-92-S1	3.4 ± 0.2	77.0	10.1	67.6	57.4	1.0	43.4	56.9	0.58	2155	7219	
TX	TX09CS1066	3.3 ± 0.2	75.3	9.7	65.3	58.2	2.4	41.3	58.5	0.60	2329	7662	
FL	FL02011-I-J2	2.4 ± 0.2	74.8	8.6	61.1	56.2	1.9	39.6	58.6	0.60	1957	4697	

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

NDF = Neutral Detergent Fiber

ADF = Acid Detergent Fiber

TDN = Total Digestable Nutrients

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

Table 20. Mean yields and agronomic characteristics of eight fall seeded oat lines evaluated for silage at the Middle Tennessee Research and Education Center for two years (2010-2011).

Brand	Variety	Dry Weight	65% Moisture			
		Avg. Yield	Avg. Yield	Moisture	(n=2)	
		± Std Err.	± Std Err.	at harvest		
(n=2)	(n=2)	(n=2)	(n=2)	(score)	inches	
TX	TAMO 406	3.5 ± 0.1	9.9 ± 0.3	67.8	1.0	31
LA	LA03063SBSBSB-S4	3.1 ± 0.1	9.0 ± 0.3	67.0	1.0	32
FL	Horizon 201	3.0 ± 0.1	8.6 ± 0.3	69.8	1.0	34
TX	TX05CS556	3.0 ± 0.1	8.5 ± 0.3	71.2	1.0	31
TX	TX07CS3697	2.9 ± 0.1	8.3 ± 0.3	70.3	1.0	33
LA	Horizon 270	2.7 ± 0.1	7.8 ± 0.3	70.5	1.0	30
TX	TX07CS2783	2.7 ± 0.1	7.8 ± 0.3	71.3	1.0	33
NC	Rodgers	2.7 ± 0.1	7.7 ± 0.3	71.0	1.0	33
Average (bu/a)		3.0	8.4	69.9	1	32
L.S.D._{.05} (bu/a)		0.4	1.1			
C.V. (%)		8.9	8.9			

Lodging = 1 to 5 scale; where 1 = 95% of plants erect; 2.5 = ~50% of plants leaning at angle ≥ 45°; 5 = 95+% of plants leaning at an angle ≥ 45°.

Table 21. Mean yields † and feed quality characteristics of eight oat varieties evaluated for silage at the Middle Tennessee Research and Education Center for two years (2010-2011).

Brand	Variety	Dry Weight	NDF 30h									Milk/ton [‡] (n=2)	Milk/acre [‡] (n=2)
		Avg. Yield	Moisture	Crude									
		± Std Err.	at Harvest	Protein	NDF	IV Digest	Starch	ADF	TDN	NEL	Mcalcs/lb	lbs/ton	lbs/acre
(n=2)	(n=2)	(n=2)	(n=2)	(n=2)	(n=2)	(n=2)	(n=2)	(n=2)	(n=2)	(n=2)			
TX	TAMO 406	3.5 ± 0.1	67.8	8.8	65.4	56.0	5.0	41.7	58.2	0.58	2331	7928	
LA	LA03063SBSBSB-S4	3.1 ± 0.1	67.0	10.0	60.4	58.9	4.6	38.5	59.3	0.59	2416	7630	
FL	Horizon 201	3.0 ± 0.1	69.8	7.9	61.0	56.9	5.7	38.7	59.1	0.59	2302	6812	
TX	TX05CS556	3.0 ± 0.1	71.2	8.8	66.1	52.5	3.1	42.9	56.7	0.56	2108	6188	
TX	TX07CS3697	2.9 ± 0.1	70.3	9.4	65.1	53.2	4.3	41.8	56.5	0.57	2177	6230	
LA	Horizon 270	2.7 ± 0.1	70.5	9.8	63.1	55.7	4.3	40.1	58.7	0.59	2381	6467	
TX	TX07CS2783	2.7 ± 0.1	71.3	10.2	60.8	57.5	4.9	38.3	60.0	0.60	2448	6648	
NC	Rodgers	2.7 ± 0.1	71.0	8.7	63.6	57.6	4.6	40.6	58.3	0.59	2390	6502	

† yields reported are dry weight basis unless otherwise indicated, feed analysis reported on an "dry weight" basis

NDF = Neutral Detergent Fiber

TDN = Total Digestable Nutrients

ADF = Acid Detergent Fiber

NEL = Net Energy for Lactation

‡ based on University of Wisconsin Milk2006 software program.

Table 22. Characteristics, as described by the seed company, of corn silage hybrids evaluated in yield tests in Tennessee during 2011.†

Brand	Hybrid §	Grain Color	Maturity	Herbicide Tolerance	BT Gene	Released or Experimental	Seed Treatment
Augusta	A5337EVT3	Y	111	RR	CB/RW	R	Poncho 250
Augusta	A5461GTCBLLA	Y	111	GT/LL	CB	R	Avicta 500
Augusta	A5462GT3000A (RR/LL/CB/RW)	Y	112	GT/LL	CB/RW	R	Avicta 500
Augusta	A5560VT3	Y	110	RR	CB/RW	R	Poncho 250
Augusta	A5464GTCBLL	Y	114	RR/LL	CB	R	Cruiser 250
Augusta	A6166GT3000A (RR/LL/CB/RW)	Y	116	RR/LL	CB/RW	R	Avicta 500
Augusta	A6867CBLL	Y	117	LL	CB	R	Cruiser 250
Augusta	A6867GTCBLLA	Y	117	RR/LL	Bt	R	Avicta 500
Augusta	A7664VT3	Y	114	RR	CB/RW	R	Poncho 250
Croplan	6286 SS (RR2/LL/CB/RW)	Y	113	RR2/LL	GCB,HXT,RW,	R	Cruiser 250
Croplan	7131VT3	Y	114	RR	CB/RW	R	Cruiser 250
Croplan	8221VT3	Y	118	RR	YGCB/RW	R	Cruiser 250
Croplan	8505VT3P	Y	117	RR	YG,CB,C,RW	R	Cruiser 250
Croplan	8756VT3	Y	118	RR	YGCB/RW	R	Cruiser 250
Croplan	9009 RH	Y	124	RR/LL	YGCB/RW	R	Cruiser 250
DeKalb	DKC67-21 GENVT3P	Y	117	RR	YG,CB,C,RW	R	Poncho 500, Votivo
Masters Choice	MC-590	Y	116	---	---	R	Poncho 250
Masters Choice	MCT-6583 (RR/LL/CB/RW)	Y	115	RR/LL	CB/RW	R	Poncho 250
Mycogen	TMF2H918 (RR/LL/HX)	Y	123	RR/LL	HX1	R	Cruiser 250
Mycogen	TMF2L872 (RR/LL/HX1)	Y	118	RR/LL	HX1	R	Cruiser 250
Wyffels	W8437 (VT3Pro)	Y	115	RR	YG,CB,C,RW	R	Metalexyl, ipconazole, trifloxystrobin, clothianidin 250
Wyffels	W8681 (VT3Pro)	Y	115	RR2	YG,CB,C,RW	R	Metalexyl, ipconazole, trifloxystrobin, clothianidin 250

Codes:

Bt, YG, YGCB, CB, HX = contains a *Bacillus thuringiensis* gene for insect resistance

CBRW, RW, CRW = contains a gene for rootworm resistance

CL = contains a gene for tolerance to Imidazolinone class herbicides

† Information on this table provided by the respective seed companies.

§ If a trait appears inside parenthesis i.e. (RR/CB), then it is not part of the hybrid name.

LL = contains a gene for tolerance to glufosinate

W = white grain

RR, R, R2, RR2, GT = contains a gene for tolerance to glyphosate

Table 23. Contact information for corn hybrid seed companies evaluated in yield tests in Tennessee during 2011.†

Company	Contact	Phone	Email	Web site	Address
Augusta Seed Corporation	Dennis Rawley Matt Rawley	540-886-6055 540-255-5902	augustaseed@aol.com		473 Tisdale Farm Ln, Stuanton, VA 24401
Croplan Genetics	Jesse Witt Keith Saum Jim Payne Ashley Plymale Matt Sowder Darrin Holder	256-221-5932 731-610-7006 901-652-0903 270-719-1570 901-355-7367 270-207-0190	jbwitt@landolakes.com kdsau@landolakes.com jpayne@ourcoop.com	www.croplangenetics.com	Consolidated Ag Products (Agrilience) and Tennessee Farmers Co-op Locations
Monsanto (Dekalb)		800-335-2676		www.monsanto.com www.dekalb.com	800 N. Lindberg Blvd, St. Louis, MO 63167
Masters Choice	Ryan Carter	618-697-8898	ryan@seedcorn.com	www.seedcorn.com	3010 St. Rt. 146 E. Anna, IL 62906
Mycogen Seed	Joe Emanuele	724-261-9115	jemanuele@dow.com	www.dowagro.com/mycogen	3563 Hilty Road, Export, PA 15632
Wyffels Hybrids Inc.	Dave Strawn	309-944-8334	dstrawn@wyffels.com	www.wyffels.com	13344 US Hwy 6, Geneseo, IL 61254

Table 24. Contact information for small grains seed companies evaluated in yield tests in Tennessee during 2010-11.

Company	Contact	Phone	Email	Web site	Address
Armor, Delta King (Cullum Seeds)	Lane Dill	901-233-0274	lanedill@jwrayseeds.com	www.cullumseeds.com	P.O. Box 178, Fisher, AR 72429
Dixie (Cache River Valley Seed)	Jason McGarrh	870-275-2779	jasonm@crvseed.com	www.crvseed.com	P.O. Box 10, Cash, AR 72421
Croplan Genetics	Jesse Witt Keith Saum Ashley Plymale	256-221-5932 731-610-7006 270-719-1570	JBWitt@landolakes.com kdsau@landolakes.com	www.croplangenetics.com	Consolidated Ag Products (Agrilience) and Tennessee Farmers Co-op Locations
	Jim Payne Matt Sowder	901-652-0903 901-355-7267	jpayne@ourcoop.com 901-355-7267	www.ourcoop.com	
Delta Grow Seed	Lee Hughes	800-530-7933	leehughes19@hotmail.com	www.deltagrow.com	P O Box 219, England, AR 72046
Dyna-Gro (Crop Production Services)	Todd Theobald	731-885-1212 765-623-1382	todd.theobald@cpsagu.com	www.dynagroseed.com	710 South First Street, Union City, TN 38621
University of Missouri	Mary Ann Quade Anne McKendry	573-884-7333 573-882-7707	quadem@missouri.edu mckendry@missouri.edu		University of MO Foundation Seed 3600 New Haven Rd Columbia, MO 65201
North Carolina State University	Paul Murphy	919-513-0000	paul_murphy@ncsu.edu		NC State University 840 Method Rd., Unit 3 Raleigh, NC 27695-7629
Ohio Seed Improvement Association	John Armstrong	614-889-1136	armstrong@ohseed.org		6150 Avery Road, Box 477 Dublin, OH 43017-0477
Pioneer Hi-Bred Int.	Michael Hughes	800-331-2475	michael.hughes@pioneer.com	www.pioneer.com	700 Boulevard South, Suite 302, Huntsville, AL 35802
Progeny	Corey Dildine	870-208-6032	corey@progenyag.com	www.progenyag.com	1529 Hwy 193, Wynne, AR 72396
Syngenta	June Hancock	870-483-7691	june.hancock@syngenta.com	www.agriprowheat.com	778 CR 680, Bay, AR 72411
Terral Seed Inc	Larry Mullen	318-231-8811	lmullen@terralseed.com	www.terralseed.com	P O Box 826, Lake Providence, LA 71254
University of Tennessee	Dennis West	865-974-8826	dwest3@utk.edu		3421 Joe Johnson Dr, Knoxville, TN 37996-4561

(continued)

Table 24. Contact information for wheat seed companies evaluated in yield tests in Tennessee during 2010-11.

Company	Contact	Phone	Email	Web site	Address
Unisouth Genetics (USG)	Stacy Burwick David Fandrich Mark Huffstetler Trey Hurt Wes Miller Billy Sellers	800-505-3133 931-967-3377 731-235-2167 731-836-7574 731-536-6251 731-538-2990	sburwick@bellsouth.net fandrichsupply@aol.com huffy1@crunet.com hurtco@bellsouth.net wes@obiongrain.com	www.usgseed.com	2640-C Nolensville Rd., Nashville, TN 37211 Fandrich Supply Co, Belvidere, TN Huffstetler & Sons Seed Inc, Greenfield, TN Hurt Seed Co. Inc, Halls, TN Obion Grain Co. Inc, Obion, TN Sellers Seed, Obion, TN
Virginia Tech	David Whitt	804-746-4884	dwhitt@vt.edu	www.virginiacrop.org	Virginia Crop Improvement Assoc. 9142 Atlee Station Rd Mechanicsville, VA 23116
Warren Seed	Lanny Warren	731-234-2921	lanny.warren@charter.net		208 South Thompson St., Union City, TN 38261
USDA (Oats)	Myron Fountain	919-513-7739	Myron.Fountain@are.usda.gov		USDA-ARS Plant Science Research Uint 1416 Gardner Hall, Dept Plant Pathology North Carolina State University, Campus Box 7616 Raleigh, NC 27695-7616